

Mixed cereal grazing trials – across site analysis 2004 – 2009, Northern NSW

Jan Edwards

District Agronomist, Cowra

James Fleming

District Agronomist, Coonabarabran

Glen Roberts

Former Research Agronomist, Temora

Frank McRae

Technical Specialist (Cereals), Orange

February 2010

Key Points

- Oats produced higher mean dry matter yields at the first dry matter assessment.
- Dry matter yields at the second assessment were similar between crop types.
- Overall oat varieties produced the largest quantity of total dry matter than other crop types.
- Triticale and barley varieties produced higher grain recovery yields than the wheat and oat varieties.
- Total dry matter was higher in the Northern trial set.
- Grain yields higher in the Southern trial set.

Trial aim

Industry and Investment NSW fund and manage a set of Mixed Cereal Grazing trials. They are sown annually at a range of locations in NSW.

The aim of the trials is to compare amounts of dry matter produced by each variety and their recovery after grazing for grain production.

Trial details

40 winter cereal varieties (3 barley varieties, 10 oats varieties, 5 triticale varieties and 22 wheat varieties) were sown at a range of locations.

The northern sites include Binnaway, Cumnock, Curban, Merrygoen, Purlewaugh, Scone, Somerton and Wongarbon. The southern sites include Cowra, Culcairn, Galong, Temora, Finley, Goulburn and Gunning. The trials were sown as small plots using a complete randomised block design with three replicates.

Trial management

The trials were sown as early as the season permitted. Seeding rates were based on recommendations in the Winter Crop Variety Sowing Guide 2009.

Fertiliser was applied according to individual site requirements. The aim was to supply enough fertiliser to achieve 4000 to 5000 kg/ha total dry matter production and 2000 to 3000 kg/ha grain yields.

Weed control varied according to individual site's needs.

Measurements

To sow, graze and harvest each variety to suit its individual maturity and vernalisation requirements is impracticable. The trial was managed to suit the majority of entries.

The three measurements taken were:

- Dry matter 1 (DM1) prior to first grazing
- Dry matter 2 (DM2) prior to second grazing
- Grain yield

Dry matter was measured using a capacitance probe and calibrated to quadrat cuts. The timing of the measurements was different at each site and determined by the season. All varieties were measured on the same day. Therefore there were slight differences in growth stage of the individual varieties.

Harvest occurred as soon as practical and when the majority of varieties reached full maturity, grain quality was not measured.

Grazing

The type and class of livestock used to graze the trials depended on the co-operator's enterprise. First grazing generally occurred from 8 weeks after sowing and completed within 7 days.

The timing of the second grazing was determined by the season. Fewer sites received a second grazing.

Trial results and discussion

The results reported are the across sites or Multi Environment Trial (MET) analysis of the northern site data from 2004-2009.

Table 1 shows the average dry matter and grain yield for the four crop types, this gives a general indication of crop performance. Please note the small number of varieties which have contributed to the barley and triticale averages.

Tables 2 to 4 present the variety data in 3 tables – dry matter 1, dry matter 2 and grain yield. Each table shows the data as an average, a percentage of the trial mean, and as a percentage of a wheat, oat or triticale standard.

Table 1: Average dry matter and grain yield for the
crop types.

NORTHE	RN NSW	Average				
		DM1	DM2	Grain yield		
Barley	3 varieties	2584	2397	2297		
Oats	10 varieties	3035	2314	1367		
Triticale	5 varieties	2498	2355	2374		
Wheat	22 varieties	2070	2240	2135		

The results show that oats produced higher average dry matter yields at the first dry matter assessment. The five highest yielding varieties were all oats – Taipan, Graza_80, Drover, Genie and Eurabbie. The first 4 were forage oat varieties.

However, by the second assessment, dry matter was similar between crop types. The five highest yielding varieties at the second assessment were Yiddah oats, Tobruk triticale, Urambie barley, Eurabbie oats and H150.2 wheat.

Overall, the oat varieties produced a larger amount of dry matter over the season than other crop types.

Triticale and barley varieties had higher grain yields than the wheat varieties. The oat varieties produced the lowest grain yields. The five highest grain yields were produced by Tobruk triticale, Urambie barley, Beaufort wheat, 98049.91 wheat and Endeavour triticale. When the two trial sets were compared, total dry matter was higher in the Northern trial set, while grain yields higher in the Southern trial set.

When selecting a suitable variety a number of factors need to be considered. The information in this publication assists with the forage and grain production quantities.

Further reading

Grain quality and disease reactions were not examined in this trial. This information, along with variety descriptions can be found in the *Winter Crop Variety Sowing Guide 2010*.

There is also a companion sheet – *Mixed cereal grazing trials* – *across site analysis* 2004 – 2009, *Southern NSW*.

Information can also be found in Primefact 720 *Cereals for grazing.*

Acknowledgements

The contributions of Steve Lyon, Dougal Pottie, Scott Boyd and I&I NSW Crop Evaluation Unit for sowing, harvesting and managing these trials are gratefully acknowledged.

Thank you to Colin Fritsch, Matthew Derrick and the staff at Temora Research Station for the dry matter assessments.

We also thank Chris Lisle (I&I NSW Wagga Wagga) for the statistical analysis of the data.

Finally, these trials could not happen without the cooperation, assistance and support of our long-term farmer co-operators.

File reference: Across sites mixed cereals 2010 N.doc

[©] State of New South Wales through Department of Industry and Investment (Industry & Investment NSW) 2010. You may copy, distribute and otherwise freely deal with this publication for any purpose, provided that you attribute Industry & Investment NSW as the owner.

Disclaimer: The information contained in this publication is based on knowledge and understanding at the time of writing (February 2010). However, because of advances in knowledge, users are reminded of the need to ensure that information upon which they rely is up to date and to check currency of the information with the appropriate officer of Industry & Investment NSW or the user's independent adviser.

NORTHERN NSW		DRY MATTER 1								
		Avera	age 1 st dry matt	er	Dry matter as a percentage of					
		kg/ha	Trial Mean %	Rank	Eurabbie Oats	Wedgetail Wheat	Endeavour Triticale	Yambla Barley		
GS5092	В	2689	112	12	85	111	102	96		
URAMBIE	В	2273	95	21	72	94	87	81		
YAMBLA	в	2791	116	9	89	115	106	100		
BIMBIL	0	2879	120	8	91	119	110	103		
DAWSON	0	3080	128	6	98	127	117	110		
DROVER	0	3221	134	3	102	133	123	115		
EURABBIE	0	3151	131	5	100	130	120	113		
GENIE	0	3167	132	4	101	131	121	113		
GRAZA80	0	3256	135	2	103	134	124	117		
MA6878W	0	3002	125	7	95	124	114	108		
MANNUS	0	2599	108	15	82	107	99	93		
TAIPAN	0	3273	136	1	104	135	125	117		
YIDDAH	0	2719	113	10	86	112	104	97		
AT573	Т	2603	108	14	83	107	99	93		
BREAKWELL	т	2564	107	16	81	106	98	92		
CRACKERJACK	т	2701	112	11	86	111	103	97		
ENDEAVOUR	т	2627	109	13	83	108	100	94		
TOBRUK	т	1993	83	32	63	82	76	71		
170	W	1764	73	40	56	73	67	63		
98049.91	w	1773	74	39	56	73	68	64		
98350.182	w	2030	84	28	64	84	77	73		
98613.86	W	1848	77	36	59	76	70	66		
AMAROK	w	2003	83	30	64	83	76	72		
BEAUFORT	W	2484	103	17	79	102	95	89		
BRENNAN	w	1998	83	31	63	82	76	72		
CS10.1010.3	w	2052	85	27	65	85	78	74		
EAGLEHAWK	w	2262	94	22	72	93	86	81		
EGA WEDGETAIL	w	2425	101	20	77	100	92	87		
FRELON	W	1887	79	34	60	78	72	68		
H150.2	W	2008	84	29	64	83	76	72		
MACKELLAR	w	1800	75	37	57	74	69	64		
MAROMBI	w	1920	80	33	61	79	73	69		
NAPAROO	w	2150	89	25	68	89	82	77		
PRESTON	w	2072	86	26	66	85	79	74		
SQP_REVENUE	Ŵ	1885	78	35	60	78	73	68		
SUN518A	Ŵ	2257	94	23	72	93	86	81		
SUN518A SUN520C	W	2237	94 93	23 24	72	93 92	85	80		
SUN520C SUN521A	W	2230 2449	93 102	24 19	78	92 101	93	88		
SUN521C	W	2449 2471	102	18	78	101	93 94	89		
TENNANT	W	1774	74	38	78 56	73	94 68	64		
	٧V		74	50	00	13	00	04		
Average No of trials		2403 29								

Table 2: Dry matter 1 (kg/ha).

NORTHERN NSW		DRY MATTER 2								
		Avera	age 2 nd dry matt	ter	Dry matter as a percentage of					
		kg/ha	Trial Mean %	Rank	Eurabbie Oats	Wedgetail Wheat	Endeavour Triticale	Yambla Barley		
GS5092	В	2385	104	7	99	104	100	102		
URAMBIE	В	2473	108	3	103	107	103	106		
YAMBLA	в	2333	102	10	97	101	98	100		
BIMBIL	0	2236	98	24	93	97	94	96		
DAWSON	0	2201	96	28	92	96	92	94		
DROVER	0	2241	98	22	93	97	94	96		
EURABBIE	о	2399	105	4	100	104	100	103		
GENIE	0	2194	96	30	91	95	92	94		
GRAZA80	0	2177	95	33	91	95	91	93		
MA6878W	0	2334	102	9	97	101	98	100		
MANNUS	0	2139	94	37	89	93	89	92		
TAIPAN	0	2222	97	26	93	97	93	95		
YIDDAH	0	2992	131	1	125	130	125	128		
AT573	Т	2329	102	11	97	101	97	100		
BREAKWELL	т	2319	102	12	97	101	97	99		
CRACKERJACK	т	2082	91	40	87	90	87	89		
ENDEAVOUR	т	2391	105	6	100	104	100	102		
TOBRUK	Т	2656	116	2	111	115	111	114		
170	W	2248	98	21	94	98	94	96		
98049.91	w	2187	96	31	91	95	91	94		
98350.182	w	2301	101	15	96	100	96	99		
98613.86	w	2269	99	20	95	99	95	97		
AMAROK	w	2239	98	23	93	97	94	96		
BEAUFORT	Ŵ	2293	100	16	96	100	96	98		
BRENNAN	Ŵ	2314	100	13	96	100	97	99		
CS10.1010.3	Ŵ	2277	101	19	95	99	95	98		
EAGLEHAWK	Ŵ	2181	95	32	91	95	91	93		
EGA_WEDGETAIL	w	2303	33 101	52 14	96	90 100	96	99		
FRELON	w	2303	100	17	90 95	99	96	98		
H150.2	w	2394	105	5	95 100	99 104	90 100	90 103		
MACKELLAR	w	2224	97	25	93	97	93	95		
MAROMBI	W	2224	93	38	89	93	93 89	95 91		
NAPAROO	W	2132	93 104	8	89 99	93 103	99	102		
	W	2378 2199	96	8 29	99 92	95	99 92	102 94		
PRESTON	W									
SQP_REVENUE	W	2145	94	36 25	89	93	90 01	92		
SUN518A		2168	95 05	35 24	90	94	91 01	93 03		
SUN520C	W	2169	95	34	90	94	91	93		
SUN521A	W	2103	92	39	88	91	88	90		
SUN521C	W	2204	96	27	92	96	92	95		
TENNANT	W	2280	100	18	95	99	95	98		
Average		2285								
No of trials		8								

Table 3: Dry matter 2 (kg/ha).

NORTHERN NSW		GRAIN RECOVERY								
		Ave	rage grain yield	d	(a percentage of	age of			
		kg/ha	Trial Mean %	Rank	Eurabbie Oats	Wedgetail Wheat	Endeavour Triticale	Yambla Barley		
GS5092	В	2235	113	11	114	106	93	104		
URAMBIE	В	2499	126	2	128	119	104	116		
YAMBLA	в	2156	109	15	110	103	90	100		
BIMBIL	0	1525	77	34	78	73	63	71		
DAWSON	0	1184	60	37	61	56	49	55		
DROVER	0	1210	61	36	62	58	50	56		
EURABBIE	0	1954	98	30	100	93	81	91		
GENIE	0	1047	53	39	54	50	44	49		
GRAZA80	0	1224	62	35	63	58	51	57		
MA6878W	0	2038	103	25	104	97	85	95		
MANNUS	0	1551	78	33	79	74	64	72		
TAIPAN	0	822	41	40	42	39	34	38		
YIDDAH	0	1117	56	38	57	53	46	52		
AT573	Т	2359	119	8	121	112	98	109		
BREAKWELL	т	2042	103	24	104	97	85	95		
CRACKERJACK	т	2400	121	6	123	114	100	111		
ENDEAVOUR	т	2405	121	5	123	114	100	112		
TOBRUK	т	2664	134	1	136	127	111	124		
170	W	2154	109	16	110	102	90	100		
98049.91	w	2419	122	4	124	115	101	112		
98350.182	w	2158	109	14	110	103	90	100		
98613.86	w	2002	101	26	102	95	83	93		
AMAROK	w	1931	97	32	99	92	80	90		
BEAUFORT	w	2486	125	3	127	118	103	115		
BRENNAN	w	1990	100	27	102	95	83	92		
CS10.1010.3	w	2083	105	21	107	99	87	97		
EAGLEHAWK	w	2130	107	17	109	101	89	99		
EGA_WEDGETAIL	w	2102	106	20	108	100	87	98		
FRELON	w	1964	99	29	100	93	82	91		
H150.2	W	2127	107	18	109	101	88	99		
MACKELLAR	w	1976	100	28	101	94	82	92		
MAROMBI	w	2370	119	7	121	113	99	110		
NAPAROO	w	2237	113	10	114	106	93	104		
PRESTON	w	2227	112	12	114	106	93	103		
SQP_REVENUE	W	2180	110	13	112	104	91	101		
SUN518A	W	2077	105	22	106	99	86	96		
SUN520C	W	2059	104	23	105	98	86	96		
SUN521A	W	2238	113	9	115	106	93	104		
SUN521C	W	2116	107	19	108	101	88	98		
TENNANT	Ŵ	1950	98	31	100	93	81	90		
Average		1985		~ '		50				
No of trials		30								

Table 4: Grain recovery (kg/ha).