



NSW DEPARTMENT OF  
PRIMARY INDUSTRIES

# **The Economic Effects of Early-Life Nutritional Constraints in Crossbred Cattle Bred on the NSW North Coast**

**Andrew R. Alford**

Livestock Research Officer, NSW Department of Primary Industries, Armidale,  
and Cooperative Research Centre for Beef Genetic Technologies

**Linda M. Cafe**

Technical Officer, NSW Department of Primary Industries, Armidale,  
and Cooperative Research Centre for Beef Genetic Technologies

**Paul L. Greenwood**

Principal Research Scientist, NSW Department of Primary Industries, Armidale,  
and Cooperative Research Centre for Beef Genetic Technologies

**Garry R. Griffith**

Principal Research Scientist, NSW Department of Primary Industries, Armidale,  
and Cooperative Research Centre for Beef Genetic Technologies

## **Economic Research Report No. 33**

May 2007



This publication is copyright. Except as permitted under the Copyright Act 1968, no part of the publication may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any way whatever without such permission.

***Abstract***

Different patterns of nutrition during pregnancy and lactation can influence cow productivity and the performance of their offspring. An experiment was conducted on the North Coast of NSW whereby “low” and “high” pasture nutritional systems were imposed on a herd of Hereford cows during pregnancy, and then again from birth to weaning, with a crossover design also imposed to select offspring with extremes of growth to birth and/or weaning. Thus, four growth groups resulted – low-low, low-high, high-low, and high-high. Piedmontese and Wagyu bulls were used. After weaning, these offspring were grown out on the NSW Northern Tablelands and then finished to heavy market weights in a feedlot. The results of the experiment indicated that restricted early-life growth resulting in differences in weight of calves at weaning persisted until slaughter at 30 months of age. Animals that were smaller at weaning remained smaller at slaughter. Some compensation occurred following restriction of growth during lactation, but not following restriction of growth during pregnancy. However, neither carcass quality nor eating quality of the beef was adversely affected by growth restriction during early-life. An economic analysis of these data was done using the Beef-N-Omics decision support package. Two methods were used. The main results showed that for the representative cattle enterprise modelled, total gross margins ranged from \$45,500 for the low-low system to \$52,600 for the high-low system. Gross margin per hectare ranged from \$114 for the low-low system to \$132 for the high-low system, while gross margin per breeding cow ranged from \$303 for the low-low system to \$387 for the high-high system. In all cases, the gross margin for those groups with foetal growth based on a higher level of nutrition exceeded their peers on a lower level of nutrition. Therefore, it is more profitable for cows and calves to have access to a high standard of nutrition during pregnancy and up to weaning than for them to have access only to a poor standard of nutrition during this time period.

***Keywords:*** beef; early-life nutrition; economic; evaluation; Australia

***JEL Code:*** Q12, Q16

**ISSN 1442-9764**

**ISBN 978 0 7347 1791 1**

***Senior Author's Contact:***

Dr Andrew Alford, NSW Department of Primary Industries Beef Industry Centre, JS Barker Building, University of New England, Armidale, NSW, 2351.

Telephone: (02) 67 701 813

Facsimile: (02) 67 701 830

Email: andrew.alford@dpi.nsw.gov.au

***Citation:***

Alford, A.R., Cafe, L.M., Greenwood, P.L. and Griffith, G.R. (2007), *The Economic Effects of Early-Life Nutritional Constraints in Crossbred Cattle Bred on the NSW North Coast*, Economic Research Report No. 33, NSW Department of Primary Industries, Armidale, May.

# Table of Contents

	<b>Page</b>
List of Tables	iv
List of Figures	v
Acknowledgements	v
Acronyms and Abbreviations Used in the Report	v
Executive Summary	vi
<b>1. Introduction</b>	<b>1</b>
<b>2. The Early-life Nutrition and Growth Experiment</b>	<b>3</b>
<b>3. Methodology for the Economic Evaluation</b>	<b>8</b>
<b>4. Results</b>	<b>15</b>
<b>5. Discussion</b>	<b>20</b>
<b>6. Conclusions</b>	<b>21</b>
<b>7. References</b>	<b>22</b>
<b>8. Appendices</b>	
<b>A. North Coast Cow-Calf Income, Costs and Herd Parameters</b>	<b>25</b>
<b>B. Northern Tablelands Growing Out/Backgrounding Costs</b>	<b>27</b>
<b>C. Feedlotting Income and Costs</b>	<b>27</b>
<b>D. Short-Term Ryegrass Gross Margin</b>	<b>28</b>
<b>E. Permanent Subtropical Pasture Gross Margin and Production Parameters</b>	<b>28</b>
<b>F. Permanent Temperate Pasture Gross Margin</b>	<b>29</b>
<b>G. Beef-N-Omics</b>	<b>30</b>

## List of Tables

	<b>Page</b>
Table 1. Calendar of management events for the entire life of the progeny from each breeding cycle	4
Table 2. Summary of growth parameters to weaning for each early-life treatment group within the 240 core progeny	5
Table 3. Summary of backgrounding growth for each early-life treatment group within the 240 core progeny	6
Table 4. Summary of feedlot intake and growth for each early-life treatment group within the 240 core progeny	6
Table 5. Summary of important carcass attributes for each early-life treatment group within the 240 core progeny	7
Table 6. Summary of feedlot growth for each early-life treatment group within the 154 progeny tested for feed intake	12
Table 7. Summary of feedlot growth for each early-life nutritional group within the 154 progeny tested for feed intake, adjusted to a feedlot entry weight of 490kg	13
Table 8. Differences in feedlot growth for each early-life nutritional group within the 154 progeny tested for feed intake, adjusted to a feedlot entry weight of 490kg, minus actual feedlot entry weight	13
Table 9. Summary of feedlot intake and growth for each early-life treatment group within the 240 core progeny, adjusted to a common feedlot entry weight (490kg in the 154 progeny)	14
Table 10. Summary of backgrounding growth for each early-life treatment group within the 240 core progeny, adjusted to a common feedlot entry weight (490kg in the 154 progeny)	14
Table 11. Gross margin budgets for each early-life nutritional treatment group, common backgrounding and feedlotting periods	16
Table 12. Sensitivity of enterprise gross margins to fed cattle price, common backgrounding and feedlotting periods	18
Table 13. Changes in the gross margin budgets for each early-life treatment group, adjusted to a common feedlot entry weight (490kg in the 154 progeny)	19

## List of Figures

	<b>Page</b>
Figure 1. Grafton pasture growth rates for weaner production phase	10
Figure 2. Glen Innes pasture growth rates for growing out/backgrounding phase	10

## Acknowledgments

The financial and in-kind support of the Cooperative Research Centre for Cattle and Beef Quality, NSW Department of Primary Industries, CSIRO Livestock Industries and the University of New England is gratefully acknowledged. We also wish to acknowledge the considerable efforts of the following research, veterinary, technical and/or farm staff: Helen Hearnshaw, David Hennessey, Lewis Molloy, Keith Newby, William Lee, Max Johnson, Eric Donoghue and Albert Martin, NSW Department of Primary Industries, Grafton; Phil Dawes, Peter Kamphorst, Peter Newman and Ross Dicker, NSW Department of Primary Industries, Glen Innes; Stuart McClelland, Joe Brunner, Bill Johns, Steve Sinclair, Reg Woodgate, Kim Quinn and Diana Perry, NSW Department of Primary Industries, Armidale; John Wilkins, NSW Department of Primary Industries, Wagga Wagga; Reid Geddes and Matt Wolcott, Beef CRC “Tullimba” Feedlot; Peter Allingham, Greg Harper and Sigrid Lehnert, CSIRO Livestock Industries, St. Lucia; John Thompson, Andrew Slack-Smith and Jason Siddell, University of New England, Armidale; Rebecca Farrell and John Bertram, Queensland Department of Primary Industries, Brisbane and Goondiwindi. We also wish to acknowledge Louise Hale who undertook an Honours project at the University of Queensland, within this research. The excellent cooperation of the management and staff of John Dee Abattoir, Warwick, in particular Warren Stiff, John Calvert and Geoff Grant, is also gratefully acknowledged. We are also grateful to James and Lyndon Mulligan of ‘Spelga Piedmontese’, the Australian Wagyu Association, and Peter Lee of ‘Waterview Wagyu’ who generously supplied the semen and the bulls used in this study. Finally, we wish to acknowledge the contribution of Dr Hutton Oddy, Meat & Livestock Australia for his role in initiating this research.

Lloyd Davies, Fiona Scott and John Mullen provided valuable comments on an earlier version of this Report.

## Acronyms and Abbreviations Used in the Report

ABARE	Australian Bureau of Agricultural and Resource Economics
ADG	Average daily gain
BEEF CRC	Cooperative Research Centre for Cattle and Beef Quality
Beef-N-Omics	A software package (see Appendix G)
CFA	Cast-for-age
CW	Carcass weight
DM	Dry matter
ME	Metabolisable energy
MSA	Meat Standards Australia
RD&E	Research, development and extension

## Executive Summary

Modern Australian beef cattle husbandry practices aim to grow cattle efficiently on pasture during the early phases of their lives, followed by the use of high quality feedstuffs during later growth phases to reduce the risk of not meeting the targeted premium market specifications. However, pasture-reliant growth of cattle is typically a prolonged process during which cattle experience widely differing nutrition levels and growth paths due to variable pasture quality and availability, climatic extremes, and constraints on management of cattle and pastures. Different patterns of nutrition during pregnancy and lactation can influence cow productivity and the performance of their offspring. This issue has assumed greater importance in recent times with the ongoing drought and the prospect of even greater variability in climate due to global warming.

An experiment was conducted on the North Coast of NSW whereby “low” and “high” pasture nutritional systems were imposed on a herd of Hereford cows during pregnancy, and then again from birth to weaning, with a crossover design also imposed to select offspring with extremes of growth to birth and/or weaning. Thus, four nutritional cow and offspring growth groups resulted – low-low, low-high, high-low, and high-high. The aim was to maximise the divergence, within animal welfare limitations, in foetal and pre-weaning growth of the progeny, and to investigate the subsequent differences in growth rates and carcass characteristics of offspring through to market weights. To cover the range of market specifications, both Piedmontese (high muscle growth and high birth weight) and Wagyu (high marbling and low birth weight) bulls were used. The progeny of the experiment were grown out or backgrounded on the NSW Northern Tablelands until about 26 months of age then finished in a feedlot for around 120 days.

The results of the experiment indicated that restricted early-life growth resulting in differences in weight of calves at weaning persisted until slaughter at 30 months of age. Animals that were smaller at weaning remained smaller at slaughter. Some compensation occurred following restriction of growth during lactation, but not following restriction of growth during pregnancy. However, neither carcass quality nor eating quality of the beef was adversely affected by growth restriction during early-life.

An economic analysis of these data was done using the Beef-N-Omics decision support package. Two different methods were used to test whether following the experimental protocols (different weights at feedlot entry) resulted in economic outcomes different from those resulting from applying typical commercial practice. The first (following the protocols) showed that for the representative cattle enterprise modelled, total gross margins ranged from \$45,500 for the low-low system to \$52,600 for the high-low system. Gross margin per hectare ranged from \$114 for the low-low system to \$132 for the high-low system, while gross margin per breeding cow ranged from \$303 for the low-low system to \$387 for the high-high system. The second method (adjusting for a common feedlot entry weight) showed slightly lower gross margins for each early-life treatment group, but the same ranking of groups. However, in all cases, the gross margin for those animals that commenced foetal growth on a high plane of nutrition and a high growth trajectory exceeded their peers on the low plane of nutrition and low growth trajectory.

It is more profitable for cows and calves to have access to a high standard of nutrition during pregnancy and up to weaning than for them to have access only to a poor standard of nutrition

during this time period. Further, if feed is in short supply and a choice has to be made, it is more profitable for cows to have access to a high standard of nutrition during pregnancy than for cows and calves to have access to a high standard of nutrition between partuition and weaning.

# 1. Introduction

Australian beef cattle producers are increasingly managing their production systems to better target specific markets for their stock. They can do this by selecting appropriate genotypes and by tailoring their nutritional management to allow better expression of genetic advantages. Market requirements range from those favoured by high yielding, lean carcasses, such as the domestic trade market, to those requiring high intramuscular fat or marbling content, such as the domestic Hotel-Restaurant-Institution market and the high end Japanese market. As a result, animals of markedly different genetic characteristics are used as terminal sires.

Modern beef cattle husbandry practices aim to grow cattle efficiently on pasture during the early phases of their lives, followed by the use of high quality feedstuffs during later growth phases to reduce the risk of not meeting the targeted market specifications. However, pasture-reliant growth of cattle is typically a prolonged process during which cattle experience widely differing nutrition levels due to variable pasture quality and availability, climatic extremes, and constraints on optimal management of cattle and pastures. Finishing cattle on grain may not be sufficient to overcome early-life nutritional deficiencies. Hence, it is important to understand the influences of different patterns of nutrition and growth during pregnancy and lactation on cows mated to sire-breeds with high muscling or marbling potential, and on the performance of their offspring. This requirement has assumed greater importance in recent times with the ongoing drought and the prospect of even greater variability in climate due to global warming.

In relation to the underlying biological relationships, foetal growth of cattle has been extensively studied in relation to calf survival. It has been shown that growth of the foetal calf can be slowed during the latter half of pregnancy by severely restricted maternal nutrition (Holland and Odde 1992). Similarly, during late pregnancy, the size of the dam can restrict the growth of foetuses with high prenatal growth potential (Ferrell 1991; Joubert and Hammond 1958). However, the consequences of foetal growth for subsequent performance, particularly in relation to carcass and eating quality characteristics at market weights, are less well understood (Greenwood *et al.* 2002, 2005) and do not appear to have been assessed in cattle differing in sire-genotype. For example, while severe growth retardation during foetal life has been shown to reduce muscle growth and increase fatness later in life in sheep (Greenwood *et al.* 1998, 2000; Villette and Theriez 1981), such retardation had lesser effects on body composition in Hereford cattle (Tudor *et al.* 1980).

Influences of pre-weaning nutrition on post-weaning growth and composition of cattle at market weights are better described (Berge 1991; Hearnshaw 1997). Nutritional restriction early in life can have prolonged effects on subsequent growth of cattle (Reardon and Everitt 1973) and severe restriction prior to weaning does not appear to result in increased carcass fatness when animals are recovered for prolonged periods on pasture (Tudor *et al.* 1980; Berge 1991; Hearnshaw 1997). However, when fed high energy diets, animals severely growth-retarded from birth to weaning were fatter at market weights than their counterparts well-grown to weaning (Tudor 1972; Tudor and O'Rourke 1980; Tudor *et al.* 1980). As with foetal growth, the influences of nutrition and growth prior to weaning on subsequent performance, and interactions between prenatal and pre-weaning nutrition and growth, in cattle of extreme sire-genotypes for muscling and marbling, have not been studied.

In the study reported here, hypotheses were proposed that cows mated to high muscle growth or high marbling potential bulls, and their offspring, will exhibit different liveweight and growth responses to divergent nutrition and growth during pregnancy and lactation. To test these hypotheses, low and high pasture quality and availability nutritional systems were imposed during pregnancy and from birth to weaning, with the aim of maximising divergence, within animal welfare limitations, in foetal and pre-weaning growth of progeny sired by Piedmontese (high muscle growth and high birth weight) and Wagyu (high marbling and low birth weight) bulls.

This study was also designed to provide subsets of animals with divergent prenatal growth (approximately 30 per cent difference in birth weight) and growth to weaning (approximately 0.5 v. 1.0 kg/day ADG) in divergent genotypes for related studies on molecular and cellular development of carcass tissues, and on carcass and eating quality characteristics. Ultimately, this information will be used to enhance models for phenotypic prediction of beef cattle performance.

Commercially, the information is also important for beef cattle producers to assess whether it is economically feasible to invest in different genetics and /or different nutrition systems.

In this report, the experimental results are reviewed, a farm-level modelling system is described that allows an economic evaluation of the experimental results, and the economic outcomes of applying this system in two different ways are reported. Implications are then drawn for beef cattle producers in the study area.

## 2. The Early-Life Nutrition and Growth Experiment

### *Overview of the Study*

This experiment was conducted as a part of the research program of the Cooperative Research Centre for Cattle and Beef Quality (the Beef CRC). The objective was to study the long-term consequences of different patterns of early-life growth on subsequent growth to heavy market weights, and on the resulting carcass yield and eating quality characteristics (Greenwood *et al.* 2005, 2006). There was also interest in comparing high yielding and high marbling types of cattle.

The dams used in the experiment were Hereford cows and heifers (360 in total), and the sires were Piedmontese (muscling) and Wagyu (marbling) (four of each). The experiment was conducted at Grafton Agricultural Research and Advisory Station, on the NSW North Coast, over two breeding cycles. Mating commenced in November 2000, with calves being born in the winter/spring of 2001 and again in the winter/spring of 2002. Following post-weaning backgrounding and finishing on the NSW Northern Tablelands, the final progeny group was slaughtered in March 2005.

Cows were managed within ‘High’ and ‘Low’ nutritional systems at Grafton from diagnosis of pregnancy to parturition. Lactating cows and their calves were similarly managed until weaning at about 8 months of age. This produced calves with high or low birth weights, and high or low growth to weaning. Half of the animals swapped nutritional treatments at birth, resulting in progeny with the following combinations of early-life nutrition and growth:

- LL = Low during pregnancy, and Low during lactation
- LH = Low during pregnancy, and High during lactation
- HL = High during pregnancy, and Low during lactation
- HH = High during pregnancy, and High during lactation

The number of progeny reaching weaning from the two breeding cycles was 534. Of these, 240 core animals were selected, representing extremes of birth weight and growth to weaning. Following weaning, these core animals were grazed on improved temperate pastures (backgrounded) at Glen Innes Agricultural Research and Advisory Station on the New England Tablelands, in steer or heifer cohort groups until they reached around 500kg. They were then grain fed for 120 days at the Tullimba research feedlot west of Armidale. They were about 30 months of age at feedlot exit, when they were slaughtered at the John Dee Abattoir near Warwick in south-east Queensland. Carcass attributes were assessed there. This core group of 240 cattle has been used in the economic analysis, as there is a complete data set relating to each of these animals.

This experimental protocol approximated a common production system in the region where weaners are bred on the NSW North Coast and then grown out on the better quality pastures of the NSW Northern Tablelands.

The calendar of management events for progeny of the two breeding cycles is shown in Table 1, and a more detailed description is provided by Cafe *et al.* (2006b).



### *Nutritional Treatments at Grafton*

The early-life High and Low nutritional treatments were pasture-based, with the use of supplementation as required to meet the experimental growth targets and to ensure animal welfare requirements were maintained within the low nutritional systems.

The Low system was based on poor quality native and naturalised pastures on unfertilised duplex soils. The target growth rate for these calves was 500-600 grams per day.

The High system was based on improved pasture species on heavy alluvial flats and higher red alluvium. Irrigated ryegrass was used to fill the winter feed gap that prevails in this region, and to ensure continued High nutrition. The target growth rate for these calves was 800-900 grams per day.

During both breeding cycles, some strategic feed supplementation was required, but this was more significant in 2002/2003 due to the very severe drought occurring in the region at the time. Lactating cows and calves were fed a pelleted, complete ration for most of the pre-weaning period as there was very little dry matter available. The supplementation allowed the research growth targets to be met, while ensuring welfare standards were maintained (Cafe *et al.* 2006b).

The growth parameters for each early-life growth group are shown in Table 2. The calves selected for the low birth weight groups were some 10kg lighter than their high birth weight counterparts. Subsequent high nutrition up until weaning significantly raised daily gain and weaning weights, but those calves on high nutrition during pregnancy (HL and HH) were always ahead of their low nutrition counterparts (LL and LH).

**Table 2. Summary of growth parameters to weaning for each early-life treatment group within the 240 core progeny**

Early-life growth treatment	n	Birth weight (kg)	Weaning weight (kg)	Pre-weaning ADG (g/d)
LL	60	28.3	139	513
LH	60	28.9	210	826
HL	59	38.1	162	596
HH	61	39.5	233	923
s.e.d.		0.76	5.1	19.3

### *Backgrounding at Glen Innes*

Three cohorts of progeny (two steer cohorts and one heifer cohort) were sent to Glen Innes to start pasture backgrounding soon after weaning. They grazed as cohort groups (n = 80 per cohort) for approximately 18 months. The pastures at Glen Innes comprise good quality temperate improved species (Fescue, Phalaris and White clover), producing high cattle weight gain over spring and summer, but little gain in the cold winter period.

Some supplementation was required at Glen Innes also during the 2002/2003 drought to ensure animals reached their target feedlot entry weights by approximately 26 months of age.

The growth of the early-life treatment groups during backgrounding is shown in Table 3. Again, those calves with high growth rate during pregnancy (HL and HH) were always ahead of their low growth counterparts (LL and LH). However, there was also some compensatory gain evident, with those calves that entered backgrounding on a slower growth path between birth and weaning (LL and HL) having faster backgrounding growth rates than their LH and HH peers.

**Table 3. Summary of backgrounding growth for each early-life treatment group within the 240 core progeny**

Early-life growth treatment	n	Weaning weight (kg)	Feedlot entry weight (kg)	Backgrounding ADG (g/d)
LL	60	139	473	602
LH	60	210	509	540
HL	59	162	511	628
HH	61	233	553	576
s.e.d.		5.1	9.0	12.3

#### *Feedlot Finishing at Tullimba*

The 240 core cattle were grain-fed at Tullimba feedlot for approximately 120 days after backgrounding. The feedlot growth and exit weights are shown in Table 4. Feedlot entry was based on cohort age, not on individual weights as would be done commercially. At the completion of feedlotting, the cattle were approximately 30 months of age. It is interesting that there is no evidence of compensatory gain in the feedlot – those animals that went in lighter grew more slowly than those that went in heavier.

**Table 4. Summary of feedlot intake and growth for each early-life treatment group within the 240 core progeny**

Group	n	Feedlot entry weight (kg)	Feed intake (t/hd/120d)	Feedlot exit weight (kg)	Feedlot ADG (g/d)
LL	60	473	1.51	629	1 465
LH	60	509	1.61	666	1 498
HL	59	511	1.68	682	1 598
HH	61	553	1.76	723	1 640
s.e.d.		9.0		13.2	61.9

#### *Carcass Characteristics*

Following feedlotting, the cattle were transported to John Dee Abattoir at Warwick in southern Queensland for slaughter and assessment of carcass attributes. Detailed carcass, yield and meat quality data were collected, a summary of which is shown in Table 5. Generally, the carcasses from all four treatments met the market specifications for the grain-fed export market that the abattoir supplies (however, examination of the P8 measures suggests that if specifications are tightened sufficiently, the HH pre weaning treatment in

particular would be penalised more often given that industry currently aim for less than 18mm of fat).

**Table 5. Summary of important carcass attributes for each early-life treatment group within the 240 core progeny**

Early-life growth treatment	n	Carcass weight (kg)	Hot P8 fat depth (mm)	Retail meat yield (kg/animal)	Retail meat yield (%)	US marble score
LL	60	353	19.6	234	67.4	456
LH	60	376	21.2	243	65.7	441
HL	59	383	19.2	250	66.4	449
HH	61	409	22.0	265	65.5	436
s.e.d.		7.6	1.4	5.7	0.56	22.7

### *Conclusion*

Restricted early-life growth resulting in differences in weight of calves at weaning persisted until 30 months of age. Animals that were smaller at weaning remained smaller at slaughter. Some compensation occurred following restriction of growth during lactation, but not following restriction of growth during pregnancy.

Carcass quality of these animals was not adversely affected by growth restriction during early-life, under present carcass specifications. Smaller animals had smaller carcasses, but tended to yield a higher proportion of meat than the larger animals. This was mainly due to the weight differences placing them at different stages of the growth curve, so that the larger animals had undergone more fattening than the smaller ones. Subcutaneous and intramuscular fat measures were not affected to the extent of influencing the value of the carcass. These findings have subsequently been confirmed in cattle grown slowly or rapidly to weaning and backgrounded to the same feedlot entry weights (Cafe *et al.* 2006a).

Similarly, eating quality of beef was unaffected by growth during early-life (see Greenwood *et al.* 2006), hence no differences in returns would occur if an objective beef marketing system such as Meat Standards Australia was used.

One aspect of the experimental design to remember when interpreting the outcomes is that animals were deliberately selected to exhibit extremes of growth from the nutritional treatments. This necessarily introduces a bias against the low groups and a bias toward the high groups. In this sense the experiment and the analyses that have followed are best-case vs worst-case scenarios.

### 3. Methodology for the Economic Evaluation

The nature of the experimental protocols resulted in a number of decisions being made that would not be consistent with normal commercial practice. For example, the very poor seasonal conditions at Grafton during much of the experimental phase necessitated the use of large levels of supplementation of some cow treatment groups to obtain the targeted high nutritional planes. These levels and consequent costs of feed supplements would be obviously uneconomic in commercial beef production terms. Further, because of the experimental protocols, feedlot entry was based on cohort age, not on individual weights as would be done commercially, and slaughter was also based on age rather than a target weight (although in this case the carcasses from all four treatments met the market specifications of the processor).

Therefore it was decided not to model the experimental data exactly as recorded, but to examine the implications of the experimental outcomes for a commercial producer by incorporating the key results into a representative cattle enterprise model. The limitations of this methodological approach to extrapolation of trial data to farm level analyses can be addressed to some extent through the appropriate validation of the model used and the use of sensitivity analyses of key assumptions (Dillon and Anderson 1990). See also the discussion in Davidson and Martin (1965) on this topic.

A farm level economic evaluation of the experimental outcomes was undertaken following the three production phases outlined above. This would be consistent with a representative farmer having a cow-calf breeding enterprise on the NSW North Coast as well as a growing-backgrounding enterprise in the adjacent Northern Tablelands region. Traditionally, North Coast beef production systems have focused upon store cattle breeding (Davies *et al.* 1999) while a broader range of enterprises such as breeding and growing out of beef cattle as well as the transfer in of store cattle from coastal regions for growing and finishing is typical of beef production in the Northern Tablelands (Alford *et al.* 2003). The farmer is also assumed to custom finish his cattle in a local feedlot.

The farmer is assumed to have 200ha of mixed pasture available on the North Coast, and another 200ha of mixed pasture available on the Northern Tablelands.

The economic evaluation is based on 2006 average prices and costs.

#### *The Beef-N-Omics Model*

The Beef-N-Omics software package (Dobos *et al.* 2006) was used in this evaluation. This package has been selected for all farm level modelling of new technologies relevant to Southern Australian beef production systems within the Beef CRC. It has been used by beef extension officers and some commercial producers over a number of years enabling practical validation of the model including the herd dynamics and metabolic energy demand algorithms derived from MAFF (1984) standards. The Beef-N-Omics program incorporates feed budgets and financial gross margin budgets for static herds. It is sufficiently flexible to allow for the input of various ages and liveweights for growing stock from weaning to turn-off.

To reduce the complexity of the economic analysis of the early-life nutritional study, it was decided to use Beef-N-Omics assuming the same land resource and associated pasture

resource for each of the growth treatments. This avoids confounding effects of different capital investment in land and associated pasture management between nutritional treatments, confining the analysis to the major focus of the study. Thus, varying areas to improved pastures was not considered. Rather, pasture types and areas for each of the two different regions were held fixed across each growth path, but energy available for the cow herd was varied by altering the stocking rate to just avoid the use of supplementary feeding and still provide sufficient metabolisable energy (ME) to meet the four sets of cattle growth rates. Thus for example, for the cow-calf phase at Grafton, 150 breeding cows could be run on the standard 200 ha of pasture to achieve the LL growth path, but only 128 breeding cows could be run on the same pasture to reach the HH growth path. Limitations of this approach are recognised given the simple ME approach used by Beef-N-Omics and the associated pasture modelling, however the methodology allows for a consistent approach across all four experimental treatments.

### *Pasture Types*

An accurate representation of pasture types and growth rates through the year is a crucial input into a Beef-N-Omics analysis. The Beef-N-Omics package contains an extensive pasture library, that has been built up from various NSW Agriculture trial data sets that were collated for use in the Prograze program (NSW Agriculture 1996). New pasture types can be added at any time.

A number of pasture growth profiles provided in the Beef-N-Omics library were tested for their similarity to the various pasture types implied by the cow/calf growth profiles for the various treatments. The different pastures mixes modelled in the representative enterprises (described below) were considered to be consistent with the pasture types and relative proportions available to the experimental treatments.

There are three pasture types assumed for the Grafton phase of the experiment:

1. Setaria (*Setaria sphacelata*), Rhodes grass (*Chloris gayana*) and White clover (*Trifolium repens*) (100ha).
2. Kikuyu (*Pennisetum clandestinum*) dominant pasture (60ha).
3. Short term ryegrass (*Lolium multiflorum*) with periodic nitrogen application (40ha).


The growth rates for these three pasture types are shown in Figure 1. The growth rates suggested for these subtropical species are consistent with those identified by Ashwood *et al.* (1992) where growth rates for Setaria and kikuyu pastures averaged between 50 and 60 kg DM/ha/day in late summer and early autumn and 5 kg/ha/day in winter.

In relation to the Glen Innes phase of the experiment, introduced pasture mixes are commonly utilised in the Northern Tablelands grazing systems to produce feeder cattle for the feedlot market or to finish beef cattle (Davies and Llewelyn 2006). Such pastures include introduced temperate grasses (eg., Phalaris [*Phalaris spp.*], Fescue [*Festuca arundinacea*], Cocksfoot [*Dactylis glomerata*], ryegrasses [*Lolium spp.*]) and legumes (eg., White clover [*Trifolium repens*] and Sub clover [*T. subterraneum*]). These pasture types and associated agronomic practices are detailed extensively for the region by various authors, for example Lodge and Whalley (1989), Lowien *et al.* (1997) and NSW Agriculture (1996).


The Beef-N-Omics model includes an introduced grass dominated pasture of Fescue (*Festuca arundinacea*) / Phalaris (*Phalaris spp.*) with at least 20 per cent base dry matter present as

White (*T. repens*) or Sub clover (*T. subterraneum*) and with annual maintenance fertilizer applications. This mix is assumed to cover the entire 200ha of available grazing land. The growth rate for this pasture mix is shown in Figure 2.

**Figure 1. Grafton pasture growth rates for weaner production phase**



**Figure 2. Glen Innes pasture growth rates for growing out/backgrounding phase**



### *Cattle Prices and Costs*

As noted above, prices and costs used in the analysis are for 2006.

Herd costs and returns for the cow-calf activity at Grafton are derived from a standard NSW Department of Primary Industries budget as shown in Appendix A. Costs for the Glen Innes backgrounding phase are shown in Appendix B. The costing of the feedlot phase of the experiment was derived from the feedlot operator. Details of feedlot costs and income for 2006 are provided in Appendix C. Apart from fixed per head charges, the cost of feedlotting is based upon feed consumed at an average price of \$260 per tonne as fed (85% DM). Combined feed intake data for each of the four treatments were used to derive average feed consumed per head in the feedlot. Average intakes (t/hd) over the 120 day feedlotting phase were 1.51, 1.61, 1.68 and 1.76 t/hd for the LL, LH, HL and HH treatments respectively. Since the carcasses from all four treatments met the market specifications for the grain-fed export market that the abattoir supplies, a common price of \$4.00/kg cw was applied to all fed cattle.

Budgets for the development and annual maintenance of the various pastures assumed in the analysis are provided in Appendix D and E for the Grafton phase, and in Appendix F for the Glen Innes phase. The cost used for the Grafton phase in the cost and return calculations is a weighted average based on the relative areas of the two pasture types. Also, the costs in the Appendices are on an annual basis – in the costs and return calculations done below, the pasture costs are each multiplied by 1.5 to account for the 18 month period that the Grafton pastures are feeding the cows and offspring to weaning, and the 18 month period that the Glen Innes pastures are feeding the backgrounding offspring to feedlot entry (Table 1).

### *Cattle Weights and Growth Rates*

Given the basic pasture, cost and return data outlined above and in the Appendices, the economic implications of the various early-life growth treatments can be derived in two different ways.

In the first approach, the farm manager is assumed to closely follow the time lines of the experiment as shown in Table 1. Thus, all animals are backgrounded for the same period of time and they all enter the feedlot at the same time but at different weights (up to 80kg difference across the treatments). They spend the same amount of time in the feedlot but grow at different rates and exit the feedlot also at different weights (up to 100kg difference). The group weights and growth rates used are those described in Tables 2-4. While this approach would not be regarded as normal commercial practice, in this particular case there were no adverse commercial implications since the carcasses from all four treatments met the market specifications for the grain-fed export market that the abattoir supplies (Table 5). Based on these assumptions, the costs and returns for each of the four growth paths, by phase and in aggregate, are shown in the first section of Chapter 4.

In the second approach, the farm manager is assumed to follow different time lines than in the experiment. The animals are backgrounded for different periods of time so as to achieve a more common feedlot entry weight. They spend the same amount of time in the feedlot, grow at more similar rates and exit the feedlot at more similar weights. This approach is more like normal commercial practice, and has been followed in other analyses of cattle experimental data (Davies *et al.* 2007) where there have been more marked feedlot exit weight differentials and substantial penalties for non-compliance with market specifications.

Since there are no actual data for this type of management for the representative farm, it has to be simulated based on assumptions. Two main issues arise:

- whether animals that are held for longer or shorter periods on backgrounding would have the same average backgrounding ADG as that measured in the experiment; and
- whether animals that enter the feedlot at heavier or lighter weights would have the same average feedlot ADG as that measured in the experiment (with the same carcass specifications).

In relation to the first issue, any longer or shorter period on backgrounding required to achieve a group weight close to the average weight of all groups (around 512 kg) would be relatively small in relation to the overall backgrounding period of 18 months (Table 1). For example, the LL group entered the feedlot at an average weight of 473 kg (Table 3). To put on the additional 39 kg at their average backgrounding ADG of 0.6kg/day would take about 65 days – or about 10 per cent of the total backgrounding period. This is a relatively minor additional period and is also at the end of the backgrounding period, so growth rates during these extra days should not be much different from the average over the whole period. We assume therefore that for any early-life treatment group the same backgrounding ADG would apply over any shorter or longer period as well.

In relation to the second issue, there is a more direct relationship between entry weight and ADG in the feedlot, and the feedlot ADGs (Table 4) are almost three times those during the backgrounding period (Table 2). We have to be more careful in our assumptions here.

Fortunately, there are some other data available on some of these animals that will assist us (Cafe, L.M., unpublished data). Of the 240 core progeny, 154 animals were subjected to a detailed feed intake analysis at the Tullimba feedlot over a 70-day period. These were the 2001 born heifer cohort and the 2002 born steer cohort. The actual recorded data for these animals are shown in Table 6. Of note is the fact that the mean entry weight of these 154 animals and the weight for each of the early-life treatment groups is less than the weight for all of the core animals reported in Table 4 above (since there are a greater proportion of heifers in this smaller subset). However, their feedlot exit weights were generally heavier than the core animals.

**Table 6. Summary of feedlot growth for each early-life treatment group within the 154 progeny tested for feed intake**

Early-life growth treatment	Feedlot entry weight (kg)	Feedlot exit weight (kg)	Feedlot ADG (g/d)	Carcass weight (kg)
LL	453	631	1.53	353
LH	480	665	1.59	374
HL	502	707	1.75	395
HH	524	733	1.76	413
s.e.d.	8.9	12.3	0.052	7.2
All	490	683	1.66	384

Statistical models were run on this data set, adjusting for a common feedlot entry weight of 490kg (the mean weight of all the 154 animals)<sup>1</sup>. The model-predicted adjusted ADG and exit weights for each treatment are shown in Table 7. As expected, the daily gain and exit weights are higher for the LL and LH groups than actually recorded, but lower for the HL and HH groups.

The differences in weights and ADG between the actual recorded data for these 154 animals and the model predictions for the common entry weight were calculated and are reported in Table 8. For the two extreme treatment groups, there are differences of over 30kg for entry weight, over 40kg for exit weight, and over 0.006 for ADG.

**Table 7. Summary of feedlot growth for each early-life nutritional group within the 154 progeny tested for feed intake, adjusted to a feedlot entry weight of 490kg**

Early-life growth treatment	Feedlot entry weight (kg)	Feedlot exit weight (kg)	Feedlot ADG (g/d)	Carcass weight (kg)
LL	490	677	1.60	379
LH	490	678	1.61	382
HL	490	692	1.73	387
HH	490	689	1.70	390
s.e.d.	na	6.9	0.059	4.4
All	490	684	1.66	384

What we would like to do now is to use this information from the 154 animals that went through the feed intake trial to predict how the 240 core animals would have grown at a common entry weight. However, since the mean weights of this group of 154 animals are different to the mean weights of the 240 core animals (490 vs 512 for entry weight, 683 vs 675 for exit weight, respectively), we cannot just apply the absolute weight differences to predict how the 240 core animals would have grown at a common entry weight. Rather, we must apply proportional weight differences, as also shown in Table 8.

**Table 8. Differences in feedlot growth for each early-life nutritional group within the 154 progeny tested for feed intake, adjusted to a feedlot entry weight of 490kg, minus actual feedlot entry weight**

Early-life growth treatment	Feedlot entry weight (kg, %)	Feedlot exit weight (kg, %)	Feedlot ADG (g/d, %)	Carcass weight (kg, %)
LL	+37, 8.2	+46, 7.3	+0.07, 4.6	+26, 7.4
LH	+10, 2.1	+13, 2.0	+0.02, 1.3	+8, 2.1
HL	-12, 2.4	-15, 2.1	-0.02, 1.1	-8, 2.0
HH	-34, 6.5	-44, 6.0	-0.06, 3.4	-23, 5.6

<sup>1</sup> These were REML analyses including effects of birth weight, pre-weaning nutrition, sex/year cohort, sire-genotype and their interactions, with feedlot entry weight as a covariate (linear, and where significant, quadratic) to predict means at equivalent feedlot entry weight.

Thus, the percentage data in Table 8 were used to adjust the feedlot weights and ADG data for the 240 core animals. These results are shown in Table 9.

**Table 9. Summary of feedlot intake and growth for each early-life treatment group within the 240 core progeny, adjusted to a common feedlot entry weight (490kg in the 154 progeny)**

Early-life growth treatment	Feedlot entry weight (kg)	Feed intake (t/hd/120d)	Feedlot exit weight (kg)	Feedlot ADG (g/d)
LL	512	1.62	658	1 573
LH	520	1.64	675	1 529
HL	499	1.65	675	1 567
HH	519	1.67	699	1 553

The calculated feedlot entry weights are not all the same, but the spread in weights, intakes and ADGs are much reduced from the raw data. The implications of these new weights for the backgrounding phase are shown in Table 10.

**Table 10. Summary of backgrounding growth for each early-life treatment group within the 240 core progeny, adjusted to a common feedlot entry weight (490kg in the 154 progeny)**

Early-life growth treatment	Feedlot entry weight (kg)	Weight difference (kg)	Background ADG (g/d)	Days difference (d)
LL	512	39	602	+65
LH	520	11	540	+20
HL	499	-12	628	-19
HH	519	-34	576	-59

The cost, revenue and profit calculations were adjusted to include the new feedlot exit weights and feed intakes, and an allowance was also made for the extra backgrounding days required by the LL and LH groups, or the fewer backgrounding days required by the LH and HH groups, to achieve the predicted “common” feedlot entry weight.

Based on these assumptions, differences in costs and returns for each of the growth paths are shown in the second section of Chapter 4.

## 4. Results

### *Common Backgrounding and Feedlotting Periods*

The gross margins for each of the four growth paths, in aggregate and per breeding cow and per hectare, are shown in Table 11 for the first approach to modelling the representative farm. The three phases are up to weaning, backgrounding, and finishing, respectively.

Under the LL scenario, where nutrition is restricted both during pregnancy and prior to weaning, 150 breeding cows can be run on the 200 ha of North Coast pasture that is described in Figure 1. Total costs over the two pasture phases are the highest under this scenario (\$90,900), since most of the costs are on a per head basis or are related to the number of breeders. Even though numbers in the feedlot are high, the costs of feedlotting are relatively low because these animals are growing more slowly and they need less feed - \$393/head over the feeding period or about \$3.28/head/day. These are the lightest animals coming out of the feedlot (353kg carcass weight) but there are 121 of them, so feedlot income is almost \$171,000. The net effect is an enterprise gross margin of \$45,500 that translates to \$114/ha or \$303/cow.

In the LH scenario, where nutrition is restricted during pregnancy but not prior to weaning, only 132 cows can be run on the standard pasture area. Since there are less cows and calves, the costs over the two pasture phases are substantially lower (\$85,200) than for the LL scenario. Feedlot costs are much the same as the LL scenario as there are fewer animals but they are growing faster and are therefore eating more - \$418/head or \$3.48/head/day. They are slightly heavier than the LL animals but there are less of them, so feedlot income is quite a bit lower at \$165,400. Overall, the enterprise gross margin is higher than the LL case by about \$1,000, which equates to \$116/ha but \$352/cow due to the lower numbers of cows.

In the case of the HL treatment, where nutrition is not restricted during pregnancy but is restricted prior to weaning, 144 breeding cows can be run on the 200 ha of pasture. Total costs over the two pasture phases (\$88,100) are second highest, and the costs of feedlotting are the highest because there are almost as many animals as in the LL scenario but they are growing more rapidly and they need more feed - \$437/head over the feeding period or about \$3.97/head/day. These animals are relatively heavy coming out of the feedlot (383kg) and there are 118 of them, so feedlot income (\$179,200) is the highest of any scenario. The overall balance is an enterprise gross margin of almost \$52,600 that translates to \$132/ha or \$365/cow.

Finally, in the HH scenario, nutrition is not restricted either during pregnancy or prior to weaning. At these consistent high energy intakes and growth rates, only 128 cows can be run on the standard pasture area. Given these low numbers of animals, the costs over the two pasture phases are the lowest of any scenario (\$84,000) and the total feedlot costs are also relatively low even though they are growing relatively faster and are therefore eating more - \$459/head or \$3.83/head/day. They are quite a bit heavier than all the other animals but there are less of them, so feedlot income is second lowest at \$170,100. On balance, the lower costs have the most impact and the enterprise gross margin is \$49,600, or at \$124/ha and \$387/cow.

**Table 11. Gross margin budgets for each early-life nutritional treatment group, common backgrounding and feedlotting periods**

Item	Value	LL			LH			HL			HH		
		Number	Cost	Income	Number	Cost	Income	Number	Cost	Income	Number	Cost	Income
Max. breeders		150											
<b>Phase 1</b>													
cfa bull	1 238	1		1 238	1		1 238	1		1 238	1		1 238
cfa cows	656	10		6 555	9		5 900	9		5 900	8		5 244
cull cows	656	17		11 144	16		10 488	17		11 143	15		9 833
Bull	1 700	2	3 400		1	1 700		1	1 700		1	1 700	
Replacement Heifers	775	31	24 025		27	20 925		30	23 250		26	20 150	
Livestock selling costs			3 607			3 231			3 451			3 068	
Health costs			2 345			2 125			2 276			2 056	
Pasture costs	88.05	200	26 415			26 415			26 415			26 415	
Cartage to backgrounding	10	126	1 260		114	1 140		122	1 220		108	1 080	
<b>Phase 2</b>													
Health costs			722			653			699			619	
Pasture costs	92.96	200	27 888			27 888			27 888			27 888	
Cartage to feedlot	10	122	1 220		111	1 110		118	1 180		105	1 050	
<b>Phase 3</b>													
Finished cattle	4.00	121 @353kg		170 852	110 @376kg		165 440	117 @383kg		179 244	104 @409kg		170 144
Feedlot costs		@\$393/hd	47 961		111@ \$418/hd	46 407		118 @ \$437/hd	51 542		105 @\$459/hd	48 157	
Induction	19.82	122	2 418		111	2 200		118	2 339		105	2 081	
Levy	5.00	122	605			550			585			520	
Cartage to abattoir	20	121	2 420		110	2 220		118	2 360		105	2 100	
<b>Total Costs and Revenues</b>			144 286	189 788	110	136 564	183 065	117	144 905	197 524	105	136 884	186 458
<b>Total GM</b>							46 501			52 619			49 574
<b>GM/ha</b>		400		45 502 113.75	400		116.25	400		131.55	400		123.93
<b>GM/cow</b>		150		303.35	132		352.28	144		365.41	128		387.30

Overall, the results of this analysis may be summarised as follows:

- For an experiment that set out to select animals that represented extremes of birth weight and growth to weaning, the economic implications are not that large. There is a 15 per cent difference in enterprise gross margin between the best case (HL) and worst case (LL) scenarios, but only an 8 per cent difference between the HL and HH groups and only a 2 per cent difference between the LL and LH groups. This provides more evidence of the relatively flat profit surfaces found in agricultural industries (Alford *et al.* 2003; Farquharson 2005; Pannell 2006).
- Given that context, it is more profitable for cows and calves to have access to a high standard of nutrition during pregnancy and up to weaning than for them to have access only to a poor standard of nutrition during this time period. The enterprise gross margin for the HH group comes to \$49,600 (\$124/ha and \$387/cow), compared to the enterprise gross margin for the LL group of \$45,500 (\$114/ha or \$303/cow). The net benefit is more than \$4,000 to the enterprise. The value of the product from the 22 extra cows able to be run on the standard 200ha of pasture does not compensate for the extra costs of backgrounding and finishing those animals and for their substantially lower carcass weights.
- Further, if feed is in short supply and a choice has to be made, it is more profitable for cows to have access to a high standard of nutrition during pregnancy than for cows and calves to have access to a high standard of nutrition between parturition and weaning. The enterprise gross margin for the HL group is \$52,600 (\$132/ha or \$365/cow), while the enterprise gross margin for the LH group is only \$46,500 (or \$116/ha or \$352/cow). The net benefit is more than \$6,000 to the enterprise, or \$16/ha.
- Overall, the HL and HH groups clearly dominate the LL and LH groups in terms of total gross margin, margin/ha and margin/cow.
- It is evident that the gross margins are the result of some complex interactions between growth rates, stocking rates, numbers of animals being fed and kilograms of beef produced. The size of the gross margins for any one early-life treatment group, and perhaps the ranking of the groups, may change with a change in the assumed value of any one of a number of influencing variables. As one example, the sensitivity of the gross margin calculations to changes in the fed cattle price is shown in Table 12. For relatively small changes away from the assumed value of \$4/kg, the LL group still exhibits the lowest gross margin, and the HL group still exhibits the highest gross margin. However, that is not the case for more extreme price changes. At very low prices, the HL group is no longer the most profitable and in fact is the second worst, while at very high prices, the LL group is no longer the least profitable.

**Table 12. Sensitivity of enterprise gross margin to fed cattle price, common backgrounding and feeding periods (\$)**

Beef price		Early-life growth treatment			
% change	\$/kg cw	LL	LH	HL	HH
-50	2.00	-39 924	-36 219	-37 003	-35 498
-25	3.00	2 789	5 141	7 808	7 038
-10	3.60	28 417	29 957	34 695	32 559
0	4.00	45 502	46 501	52 619	49 574
10	4.40	62 587	63 045	70 544	66 588
25	5.00	88 215	87 861	97 430	92 110
50	6.00	130 928	129 221	142 241	134 646

*Common Feedlot Entry Weights*

The gross margins for each of the four growth paths, in aggregate and per breeding cow and per hectare, were re-calculated for the second approach to modelling the representative farm. In particular, the cost, revenue and profit calculations were adjusted to include the new feedlot exit weights and feed intakes (Table 9), and an allowance for the extra backgrounding days required by the LL and LH groups, or the fewer backgrounding days required by the LH and HH groups, to achieve a predicted “common” feedlot entry weight (Table 10). The *changes* in the major values compared to Table 11 are given in Table 13.

- The obvious result here is that all the changes in gross margins due to the second method of calculation are negative. Thus, given the particular set of experimental data analysed here, it costs more to manage different groups of animals growing at different rates to achieve common feedlot entry weights than to manage animals to enter the feedlot at the same time. The fact that all the animals were able to meet market specifications, even though they were quite different final weights, plays a large role in this outcome. If there would have been a large number of animals discounted for not reaching specification, the impact of the second method would have been quite different.
- However, the differences in gross margins between the two methods are very small in all cases. The largest difference is for the HH group, and it is only 3.5 per cent of the base result reported in Table 11.
- Finally, the groups are ranked in the same order as for the first method – the HL group is the most profitable, followed by the HH group, the LH group and then the LL group. Thus whichever method is used, the same implications for producers would hold.

**Table 13. Changes in the gross margin budgets for each early-life treatment group, adjusted to a common feedlot entry weight (490kg in the 154 progeny)**

Item	Value	LL			LH			HL			HH		
		Number	Cost	Income	Number	Cost	Income	Number	Cost	Income	Number	Cost	Income
Max. breeders		150			132			144			128		
Changes in backgrounding pasture costs	0.50/hd/d	122@65d	3 965		111@20d	1 110		118@-19d	-1 121		105@-59d	-3 098	
Changes in finished cattle revenue	4.00/kg	121 @15kg		7 260	110 @2kg			117 @-5kg		-2 340	104 @-18kg		-7 488
Changes in feedlot costs	260/t	122 @\$28/hd	3 416		111@ \$8/hd	888	880	118 @ \$-8/hd	-944		105 @ \$-25/hd	-2 625	
Changes in total costs and revenues			7 381	7 260		1 988	880		-2 065	-2 340		-5 723	-7 488
Changes in total GM							-1 118			-275			-1 765
Changes in GM/ha		400		-121 -0.30	400		-2.80	400		-0.69	400		-4.41
Changes in GM/cow				-0.81	132		-8.47	144		-1.91	128		-13.79

\* the allowance of \$0.50/head/day for changes in pasture backgrounding cost is based on current adjustment rates (Davies et al 2007).

150

\* the new carcase weight was calculated by applying a common dressing percentage of 0.56.

## 5. Discussion

The experiment analysed here was conducted on the North Coast of NSW. “Low” and “high” pasture nutritional systems were imposed on a herd of Hereford cows during pregnancy, and then again from birth to weaning, with a crossover design also imposed to select offspring with extremes of growth to birth and/or weaning. Thus, the analyses are essentially best-case and worst-case scenarios due to this selection of extremes.

The main result shows this divergence. It is more profitable for cows and calves to have access to a high standard of nutrition during pregnancy and up to weaning than for them to have access only to a poor standard of nutrition during this time period. The enterprise gross margin for the HH group comes to \$49,600 (\$124/ha and \$387/cow), compared to the enterprise gross margin for the LL group of \$45,500 (\$114/ha or \$303/cow). The net benefit is more than \$4,000 to the enterprise, \$10/ha or \$84/cow. The value of the product from the 22 extra cows able to be run on the standard 200ha of pasture does not compensate for the extra costs of backgrounding and finishing those animals and for their lower carcass weights.

Calculating costs, returns and profits by a different method gave roughly the same numerical results, and in both cases the groups are ranked in the same order – the HL group is the most profitable, followed by the HH group, the LH group and then the LL group.

However, another result is that the economic implications are not that large. There is a 15 per cent difference in enterprise gross margin between the best case (HL) and worst case (LL) scenarios, but only an 8 per cent difference between the HL and HH groups and only a 2 per cent difference between the LL and LH groups. Given that extremes were selected, this provides more evidence of the relatively flat profit surfaces found in many agricultural industries (Alford *et al.* 2003; Farquharson 2005; Pannell 2006).

Commercially, the information is also important for beef cattle producers to assess whether it is economically feasible to invest in different genetics and /or different nutrition systems. For example, given the calculated net benefit of \$10/ha or \$84/cow between the LL and HH groups, this is the upper bound on the amount of money the typical producer could invest each year in pasture improvement to shift them from a LL growth path to a HH growth path.

It is also planned that this type of information will be used to enhance models for phenotypic prediction of beef cattle performance. This is the subject of current work in the Beef CRC.

One outcome that was not measured in this experiment was the impact of maternal nutritional setbacks on subsequent reproductive performance. It would be expected that cows that suffered a nutritional setback would take longer to re-conceive and/or exhibit a lower re-conception rate than cows that had a high standard of nutrition during their previous pregnancy or prior to re-conception. There is no Australian evidence on this issue, but there is some recent work in the US (Stalker *et al.* 2006).

We have not discussed sire-breed effects, although these aspects of the overall experimental design are covered in Cafe *et al.* (2006b) and in Greenwood *et al.* (2006). In the core data set, sire breed and early-life nutrition or growth interactions were not evident for the commercial characteristics of interest.

## 6. Conclusions

Different patterns of nutrition during pregnancy and lactation can influence cow productivity and the performance of their offspring. An experiment was conducted on the North Coast of NSW whereby “low” and “high” pasture nutritional systems were imposed on a herd of Hereford cows during pregnancy, and then again from birth to weaning, with a crossover design also imposed. Thus, four nutritional groups resulted – low-low, low-high, high-low, and high-high. Piedmontese and Wagyu bulls were used. After weaning, the offspring of these cows were grown out on the NSW Northern Tablelands and then finished to heavy market weights in a feedlot.

The results of the experiment indicated that restricted early-life growth resulting in differences in weight of calves at weaning persisted until 30 months of age. Animals that were smaller at weaning remained smaller at slaughter. Thus the hypotheses proposed “that cows mated to high muscle growth or high marbling potential bulls, and their offspring, will exhibit different liveweight and growth responses to divergent nutrition and growth during pregnancy and lactation” could not be rejected. Some compensation occurred following restriction of growth during lactation, but not following restriction of growth during pregnancy. However, neither carcass quality nor eating quality of the beef was adversely affected by growth restriction during early-life.

An economic analysis of these data was undertaken using the Beef-N-Omics decision support package. Two different methods were used. The main method showed that for the representative cattle enterprise modelled, total gross margins ranged from \$45,500 for the low-low system to \$52,600 for the high-low system. Gross margin per hectare ranged from \$114 for the low-low system to \$132 for the high-low system, while gross margin per breeding cow ranged from \$303 for the low-low system to \$387 for the high-high system. In all cases, the gross margin for those groups that started life on a higher level of nutrition exceeded their peers on a lower level of nutrition.

A slightly different type of analysis showed that it costs more to manage different groups of animals growing at different rates to achieve common feedlot entry weights than to manage animals to enter the feedlot at the same time. However, the differences in gross margins between the two methods are small in all cases. The largest difference is for the HH group, and it is only 3.5 per cent of the base result reported in Table 11. And in both methods the groups are ranked in the same order – the HL group is the most profitable, followed by the HH group, the LH group and then the LL group. Thus whichever method is used, the same overall findings are robust and the same implications for producers would hold.

While the economic implications are not that large, it is more profitable for cows and calves to have access to a high standard of nutrition during pregnancy and up to weaning than for them to have access only to a poor standard of nutrition during this time period. Further, if feed is in short supply and a choice has to be made, it is more profitable for cows to have access to a high standard of nutrition during pregnancy than for cows and calves to have access to a high standard of nutrition between parturition and weaning.

## 7. References

- Alford, A.R., Griffith, G.R. and Davies, B.L. (2003), *Livestock Farming Systems in the Northern Tablelands of NSW: An Economic Analysis*, Economic Research Report No 12, NSW Agriculture, Orange.
- Ashwood, A., Griffiths, C. and Moore, K. (1992), *Milkonomics – Feedplan*, NSW Agriculture, Wollongbar.
- Berge, P. (1991), “Long-term effects of feeding during calthood on subsequent performance of beef cattle (a review)”, *Livestock Production Science* **28**, 179-201.
- Cafe, L.M., Hearnshaw, H., Hennessy, D.W. and Greenwood, P.L. (2006a), “Growth and carcass characteristics at heavy market weights of Wagyu-sired steers following slow or rapid growth to weaning”, *Australian Journal of Experimental Agriculture* **46**, 951-955.
- Cafe, L.M., Hennessy, D.W., Hearnshaw, H., Morris, S.G. and Greenwood, P.L. (2006b), “Influences of nutrition during pregnancy and lactation on birth weights and growth to weaning of calves sired by Piedmontese or Wagyu bulls”, *Australian Journal of Experimental Agriculture* **46**, 245-255.
- Davidson, B.R. and Martin, B.R. (1965), “The relationship between yields on farms and in experiments”, *Australian Journal of Agricultural Economics* **9**(2), 129-140.
- Davies, L., Wilkins, J., Arnott, R. and Rayner, A. (1999), *Review of the store cattle industry in NSW 1998*, Animal Industries Report 7, NSW Agriculture, Orange.
- Davies, L. and Llewelyn, D. (2006), *Recent beef production and management trends in NSW*, NSW Department of Primary Industries, Orange. Available at: <http://www.agric.nsw.gov.au/reader/beefbudinfo/beef-bud-info-systems-trends.htm>
- Davies, L., McKiernan, W., Wilkins, J. and Griffith, G. (2007), “Methodology to determine gross margins in the stage prior to feedlot entry”, unpublished paper, CRC for Beef Genetic Technologies, Armidale, January.
- Dillon, J.L. and Anderson, J.R. (1990), *The Analysis of Response in Crop and Livestock Production*, Third edition, Pergamon Press, Oxford.
- Dobos, R., Carberry, P. and Davies, L. (2006), *Beef-N-Omics for Windows, User’s Manual*, NSW Department of Primary Industries, Orange, July.
- Ferrell, C.L. (1991), “Maternal and fetal influences on uterine and conceptus development in the cow: I. Growth of the tissues of the gravid uterus”, *Journal of Animal Science* **69**, 1945-1953.
- Farquharson, Bob (2006), “Production response and input demand in decision making: nitrogen fertilizer and wheat growers”, *Australasian Agribusiness Review*, paper 7.

Greenwood, P.L., Hunt, A.S., Hermanson, J.W. and Bell, A.W. (1998), “Effects of birth weight and postnatal nutrition on neonatal sheep: I. Body growth and composition, and some aspects of energetic efficiency”, *Journal of Animal Science* **76**, 2354-2367.

Greenwood, P.L., Hunt, A.S., Hermanson, J.W. and Bell, A.W. (2000), “Effects of birth weight and postnatal nutrition on neonatal sheep: II. Skeletal muscle growth and development”, *Journal of Animal Science* **78**, 50-61.

Greenwood, P.L., Wolcott, M., Hearnshaw, H., Hennessy, D.W., Morris, S.G. and Harper, G. (2002), “Fetal growth capacity influences nutritional status of Hereford cows during pregnancy”, *Animal Production in Australia* **24**, 304.

Greenwood, P.L., Cafe, L.M., Hearnshaw, H. and Hennessy, D.W. (2005), “Consequences of nutrition and growth retardation early in life for growth and composition of cattle and eating quality of beef”, *Recent Advances in Animal Nutrition Australia* **15**, 183-195.

Greenwood, P.L., Cafe, L., Hearnshaw, H., Hennessy, D.W. and Morris, S.G. (2006), “Long-term consequences of birth weight and growth to weaning for carcass, yield and beef quality characteristics of Piedmontese- and Wagyu-sired cattle”, *Australian Journal of Experimental Agriculture* **46**, 257-269.

Hearnshaw, H. (1997), “Effect of pre-weaning nutrition on post-weaning growth carcass and meat quality traits”, In: *Growth and Development of Cattle. Proceedings of the Growth and Development Workshop*, pp. 59–67. (eds. D.W. Hennessy, S.R. McLennan and V.H. Oddy), Cooperative Research Centre for Cattle and Beef Quality, Armidale.

Holland, M.D. and Odde, K.G. (1992), “Factors affecting calf birth weight: A review”, *Theriogenology* **38**, 769-798.

Joubert, D.M. and Hammond, J. (1958), “A cross-breeding experiment with cattle, with special reference to maternal effect in South Devon-Dexter crosses”, *Journal of Agricultural Science, Cambridge* **51**, 325-341.

Lodge, G.M. and Whalley, R.D.B. (1989), *Native and Natural Pastures on the Northern Slopes and Tablelands of NSW: A Review and Annotated Bibliography*, Technical Bulletin 35, NSW Agriculture, Sydney.

Lowien, J., Duncan, M., Collett, I. and McDonald, W. (1997), *A Guide to Better Pastures on the Northern Tablelands*, NSW Agriculture, Orange.

MAFF (1984), *Energy Allowances and Feeding Systems for Ruminants*, U.K. Ministry of Agriculture, Fisheries and Food, Technical Bulletin No. 433, HMSO, London.

NSW Agriculture (1996), *Prograze: profitable, sustainable grazing*, NSW Agriculture, Orange.

NSW DPI (2006), Beef gross margins budgets for 2006.  
Available <http://www.agric.nsw.gov.au/reader/beefbud>

NSW DPI (2006), Crop and pasture budgets for 2006.

Available <http://www.agric.nsw.gov.au/reader/wincropbud>

NSW DPI (2006), Guide to tractor and implement costs.

Available <http://www.agric.nsw.gov.au/reader/machine+water>

Pannell, David (2006), “Flat earth economics: the far-reaching consequences of flat payoff functions in economic decision making”, *Review of Agricultural Economics* **28**(4), 553-566.

Reardon, T.F. and Everitt, G.C. (1973), “Effects of preweaning nutrition on subsequent growth rate, feed conversion efficiency and carcass composition of identical twin steers”, *Proceedings of the New Zealand Society of Animal Production* **33**, 26-38.

Stalker, L.A., Adams, D.C., Klopfenstein, T.J., Feuz, D.M. and Funston, R.N. (2006), “Effects of pre- and postpartum nutrition on reproduction in spring calving cows and calf feedlot performance”, *Journal of Animal Science* **84**, 2582-2589.

Tudor, G.D. (1972), “The effect of pre- and post-natal nutrition on the growth of beef cattle. I. The effect of nutrition and parity of the dam on calf birth weight”, *Australian Journal of Agricultural Research* **23**, 389-395.

Tudor, G.D. and O’Rourke, P.K. (1980), “The effect of pre- and post-natal nutrition on the growth of beef cattle. II. The effect of severe restriction in early postnatal life on growth and feed efficiency during recovery”, *Australian Journal of Agricultural Research* **31**, 179-189.

Tudor, G.D., Utting, D.W. and O’Rourke, P.K. (1980), “The effect of pre- and post-natal nutrition on the growth of beef cattle. III. The effect of severe restriction in early postnatal life on the development of the body components and chemical composition”, *Australian Journal of Agricultural Research* **31**, 191-204.

Villette, Y. and Theriez, M. (1981), “Effect of birth weight on lamb performances. II. Carcass and chemical composition of lambs slaughtered at the same weight”, *Annales de Zootechnie* **30**, 169-182.

## Appendix A: North Coast Cow-Calf Income, Costs and Herd Parameters

Enterprise Unit: 100 cows  
Representative Year - 2006 dollar values

<b>Income:</b>						<b>\$</b>
42	Steer weaners	<i>To backgrounding/feedlotting</i>				
42	Heifer weaners	<i>To backgrounding/feedlotting</i>				
6	CFA cows	230 kg d.w. \hd	@ 285 ¢/kg d.w.		3 933.00	
12	Cull cows	230 kg d.w. \hd	@ 285 ¢/kg d.w.		7 866.00	
1	Cull bulls	450 kg d.w. \hd	@ 275 ¢/kg d.w.		1 237.50	
<b>Total Income</b>					<b>\$13 036.50</b>	
<b>Variable Costs:</b>						
Animal health -	vaccination, drenching and vet costs				<b>\$</b>	
	Cows	100	@ \$12.80	\hd	1 280.00	
	Bulls	3	@ \$98.33	\hd	294.99	
	Calves	84	@ \$2.43	\hd	204.12	
Ear tags	Heifers	20	@ \$15.68	\hd	313.60	
<b>Selling Costs</b>						
Cartage	Sales/ Purchases	39	@ \$7.00	\hd	273.00	
Commission	Sales Revenue	\$ 13 036.50	@ 3.5%		456.28	
Yard dues	No. of head	19	@ \$3.00	\hd	57.00	
MLA levy	No. of head	19	@ \$5.00	\hd	95.00	
Tail tags	No. of head	19	@ \$0.11	\hd	2.09	
NLIS tags	No. of head	84	@ \$2.90	\hd	243.60	
Freight to abattoir	kg dw.	4 590	@ \$0.05	\kg dw	229.50	
Cartage to backgrounding	No. of head	84	@ \$10.00	\hd	840.00	
<b>Replacements</b>						
	- Heifers	20	@ \$775.00	\hd	15 500.00	
	- Bull	1	@ \$1 700.00	\hd	1 700.00	
<b>Total Costs</b>					<b>\$21 489.18</b>	

Various prices taken from NSW DPI (2006) Beef gross margins budgets for 2006 (including herd health costs) for north coast herds. Available <http://www.agric.nsw.gov.au/reader/beefbud> (Viewed 6 Dec 2006).

---

**Herd parameters:**

---

Calving date	July - September
Weaning rate	84 %
Average weaning age	7 months
Adult mortality	2 %
Yearling mortality	2 %
Calf mortality	2 %
Bull requirement	3 %
Bull cull rate	25 %
Heifers first joined	2 years
Cow age at last joining	9 years
Other culls	5 %

---

**Age structure**

---

<b>Cow age (years)</b>	<b>Number</b>
2	20
3	17
4	15
5	13
6	11
7	9
8	8
9	7
Total joined	100
cfa cows	6

---

## Appendix B: Northern Tablelands Growing Out (Backgrounding) Costs

(based on the base North Coast 100 cow breeding herd)

### Variable Costs

Animal health -	vaccination, drenching and vet costs					\$
	Yearlings	86	@	\$5.73	\hd	492.78
Cartage to feedlot	Sales/ Purchases	84	@	\$10.00	\hd	840.00
<b>Total Costs</b>						<b>\$1 332.78</b>

Animal health costs taken from NSW DPI (2006) Beef gross margins budgets for 2006. Available <http://www.agric.nsw.gov.au/reader/beefbud> (Viewed 6 Dec 2006).

## Appendix C: Feedlotting Income and Costs

(based on the base North Coast 100 cow breeding herd)

<b>Income:</b>						\$
42	Steers	<i>Refer to Table 5 for final weights</i>	@	\$4.00/kg cw <sup>1</sup>		<i>Table 11</i>
42	Heifers	<i>Refer to Table 5 for final weights</i>	@	\$4.00/kg cw <sup>1</sup>		<i>Table 11</i>

### Variable Costs<sup>2</sup>:

	vaccination, drenching and vet costs					\$
Animal induction & Shire levy	No. of head	84	@	\$19.82	\hd	1 664.88
Feedlotting costs	Tonnes fed \hd	84	@	\$260.00	\t as fed	
Selling Costs						
MLA levy	No. of head	84	@	\$5.00	\hd	420.00
Cartage to abattoir	No. of head	84	@	\$20.00	\hd	1 680.00

<sup>1</sup> Prices for feedlot cattle are not available publicly. However the National Livestock Reporting Service does quote a price for southern Queensland 100 day grainfed cattle 300-420 cwt kg, dentition 2-4. As at 19/1/07 this category had a price range of 345 to 385 c/kg cwt (average of 366 c/kg cwt). Given that over 2006 the EYCI averaged 341.8 c/kg cwt, and at 19/1/07 the indicator was 305.5 a difference of 36.3 c/ kg cwt (or 10% lower) than the 2006 average, then a value of 400 c/kg cwt is a reasonable estimate of feedlot finished stock as described for the propose of this report.

<sup>2</sup>Feedlot costs provided by commercial feedlot reflecting the average cost per head during 2006.

## Appendix D: Short-term ryegrass gross margin

Enterprise Unit: hectare

Representative Year – 2006 dollar values

Assumes seed is spread with superphosphate application and then irrigated, after being appropriately heavily grazed.

Variable costs	Rate	Unit		Unit	TOTAL (\$/ha)
Seed					
Ryegrass (Concord)	40	kg/ha	@	\$4.00 per kg	160.00
Fertiliser					
SuperP (spread)	100	kg/ha	@	\$420 per tonne	42.00
Urea (spread)	125	kg/ha	@	\$670 per tonne	83.75
Water					
Pumping costs (over 7 appl.)	2	ML/ha	@	\$30.00 per ML	60.00
<b>TOTAL ANNUAL COSTS</b>					<b>\$ 345.75 /ha</b>

\* Fertiliser and seed costs from NSW DPI obtained from NSW DPI crop and pasture budgets for 2006. Available <http://www.agric.nsw.gov.au/reader/wincropbud> (Viewed 6 Dec 2006) or direct from produce supplier.

## Appendix E: Permanent Subtropical Pasture gross margin and production parameters

Setaria (*Setaria sphacelata*), Rhodes grass (*Chloris gayana*) and White clover (*Trifolium repens*)

This is a permanent pasture which relies upon the clover seedbank to maintain legume component therefore inputs are minimal with only maintenance fertilizer of superphosphate required.

Enterprise Unit: hectare

Representative Year – 2006 dollar values

Variable costs	Rate	Unit		Unit	TOTAL (\$)
Maintenance Fertiliser					
SuperP (spread)	150	kg/ha	@	\$420 per tonne	63.00/ha

\* Fertiliser and seed costs from NSW DPI obtained from NSW DPI crop and pasture budgets for 2006. Available <http://www.agric.nsw.gov.au/reader/wincropbud> (Viewed 6 Dec 2006) or direct from produce supplier.

## Appendix F: Permanent Temperate Pasture Gross Margin

Enterprise Unit: hectare

Representative Year – 2006 dollar values

	Rate	Unit		Unit	TOTAL (\$/ha)
Establishment costs <sup>1,2</sup>					
Disc (x2)	0.58 x2	hrs/ha	@	\$19.32 per hour	22.41
Scarifier (x2)	0.42 x2	hrs/ha	@	\$17.22 per hour	14.46
Sowing - combine	0.29	hrs/ha	@	\$25.49 per hour	7.39
Seed					
Demeter Fescue	5	kg/ha	@	\$5.00 \kg	25.00
Australian Phalaris	2.5	kg/ha	@	\$8.70 \kg	21.75
Perennial ryegrass (Impact)	1.5	kg/ha	@	\$5.50 \kg	8.25
White Clover (Haifa)+ innoc, lime	1	kg/ha	@	\$4.50 \kg	4.50
Fertiliser					
Starter	125	kg/ha	@	\$530 per tonne	66.25
Herbicides					
Glyphosate 450 (x2)	1.2 x2	L/Ha	@	\$5.00 per litre	12.00
Spray application (x2)	0.1 x2	Hrs/ha	@	\$22.16 per hour	4.43
<b>TOTAL ESTABLISHMENT COSTS</b>					<b>\$ 186.44</b>
<b>Variable costs</b>	<b>Rate</b>	<b>Unit</b>		<b>Unit</b>	<b>TOTAL (\$)</b>
Lifespan of Perennial Pasture				15 years	
<b>ESTABLISHMENT COSTS PER YEAR – Amortized*</b>					<b>17.96</b>
Maintenance Fertiliser					
Years 2-4 SuperP (spread)	375	kg/ha	@	\$420 per tonne	157.50
Years 5-15 SuperP (spread)	125	kg/ha	@	\$420 per tonne	52.50
				Average annual maintenance fertiliser	75.00
<b>TOTAL ANNUAL COSTS</b>					<b>\$ 92.96 /ha</b>

\* Pasture establishment costs were amortized using a 5 per cent interest rate and 15 year term.

<sup>1</sup>Tractor and implement variable costs from DPI Guide to Tractor and Implement Costs for a 57 KW (77HP) PTO tractor. Available <http://www.agric.nsw.gov.au/reader/machine+water> (Viewed 6 Dec 2006)

<sup>2</sup> Fertiliser and seed costs from NSW DPI obtained from NSW DPI crop and pasture budgets for 2006. Available <http://www.agric.nsw.gov.au/reader/wincropbud> (Viewed 6 Dec 2006) or direct from produce supplier.

## Appendix G: Beef-N-Omics

The Beef-N-Omics computer package (Dobos *et al.* 2006) is designed to analyse the effects that different management practices have on the profitability of a beef herd. The program integrates herd structures, feed budgets and financial gross margin budgets for beef cattle breeding herds.

User inputs are required on aspects of the beef enterprise such as herd size, live weight, calving times, age and weight at turn off, market prices, seasonal pasture growth, and variable costs. The package calculates gross margin per cow, per \$100 capital, per hectare and per tonne dry matter (DM), as well as the monthly feed surplus or deficit.

Adjustments to herd size, monthly pasture growth, months of calving, age and weight of turn off, sale prices, variable costs, cow size, weaning percentage, or other aspects of herd management can be made to assess their impact on feed requirements and subsequently on herd gross margins. Adjustments to any of those parameters will be reflected in changes in monthly feed consumption and herd gross margin from which the principles of beef cattle management can be reinforced.

Beef-N-Omics is a static herd model designed so that all the inputs are used in the calculations. This assumes that these inputs have been the same for the entire history of the herd being analysed.

Because of this, Beef-N-Omics cannot be used accurately to assess the outcome of changes to aspects like sales policy, breeding or culling policy or calving patterns which will only be applied for a year or two, for example, during droughts.

Beef-N-Omics is not a FULL biological model. Local estimates can be used, but if accurate information is available, then more precise reports are generated. A disadvantage with this approach is that users must remember to input all the correlated consequences of any change to major inputs. A misleading output could result if this is not the case.

Examples are provided in the User's Manual.

# NSW Department of Primary Industries

## Economic Research Report Series

(All available at <http://www.agric.nsw.gov.au/reader/10550>)

### Number

- 1 Brennan, J.P. and Bantilan, M.C.S. 1999, *Impact of ICRISAT Research on Australian Agriculture*, Report prepared for Australian Centre for International Agricultural Research, Economic Research Report No. 1, NSW Agriculture, Wagga Wagga.
- 2 Davies, B.L., Alford, A. and Hollis, G. 1999, *Analysis of ABARE Dairy Data for Six Regions in NSW 1991-92 to 1996-97*, Economic Research Report No 2, NSW Agriculture, C.B. Alexander College, Paterson.
- 3 Brennan, J.P. and Singh, R.P. 2000, *Economic Assessment of Improving Nutritional Characteristics of Feed Grains*, Report prepared for Grains Research and Development Corporation, Economic Research Report No. 3, Wagga Wagga.
- 4 Zhao, X., Mullen, J.D., Griffith, G.R., Griffiths, W.E. and Piggott, R.R. 2000, *An Equilibrium Displacement Model of the Australian Beef Industry*, Economic Research Report No 4, NSW Agriculture, Armidale.
- 5 Griffith, G., F'Anson, K., Hill, D., Lubett, R. and Vere, D. 2001. *Previous Demand Elasticity Estimates for Australian Meat Products*, Economic Research Report No 5, NSW Agriculture, Armidale.
- 6 Griffith, G., F'Anson, K., Hill, D. and Vere, D. 2001. *Previous Supply Elasticity Estimates for Australian Broadacre Agriculture*, Economic Research Report No 6, NSW Agriculture, Armidale.
- 7 Patton, D.A. and Mullen, J.D. 2001, *Farming Systems in the Central West of NSW: An Economic Analysis*, Economic Research Report No. 7, NSW Agriculture, Trangie.
- 8 Brennan, J.P. and Bialowas, A. 2001, *Changes in Characteristics of NSW Wheat Varieties, 1965-1997*, Economic Research Report No. 8, NSW Agriculture, Wagga Wagga.
- 9 Mullen, J.D. 2001, *An Economic Perspective on Land Degradation Issues*, Economic Research Report No. 9, NSW Agriculture, Orange.
- 10 Singh, R.P., Mullen, J.D., and Jayasuriya, R. 2005, *Farming Systems in the Murrumbidgee Irrigation Area in NSW: An Economic Analysis*, Economic Research Report No. 10, NSW Department of Primary Industries, Yanco.
- 11 Brennan, J.P., Aw-Hassan, A., Quade, K.J. and Nordblom, T.L. 2002, *Impact of ICARDA Research on Australian Agriculture*, Economic Research Report No. 11, NSW Agriculture, Wagga Wagga.
- 12 Alford, A., Griffith, G. and Davies, L. 2003, *Livestock Farming Systems in the Northern Tablelands of NSW: An Economic Analysis*, Economic Research Report No. 12, NSW Agriculture, Armidale.
- 13 Alford, A., Griffith, G. and Cacho, O. 2003, *A Northern Tablelands Whole-Farm Linear Program for Economic Evaluation of New Technologies at the Farm Level*, Economic Research Report No. 13, NSW Agriculture, Armidale.
- 14 Mullen, J.D. and Vere, D.T. 2003, *Research and Extension Capabilities - Program Economists in New South Wales Agriculture*, Economic Research Report No. 14, NSW Agriculture, Orange.
- 15 Farquharson, R.J., Griffith, G.R., Barwick, S.A., Banks, R.G. and Holmes, W.E. 2003, *Estimating the Returns from Past Investment into Beef Cattle Genetic Technologies in Australia*, Economic Research Report No. 15, NSW Agriculture, Armidale.

- 16 O'Donnell, C.J., Griffith, G.R., Nightingale, J.J. and Piggott, R.R. 2004, *Testing for Market Power in Multiple-Input, Multiple-Output Industries: The Australian Grains and Oilseeds Industries*, Technical Report for the Rural Industries Research and Development Corporation, Economic Research Report No. 16, NSW Agriculture, Armidale.
- 17 Brennan, J.P., Martin, P.J. and Mullen, J.D. 2004, *An Assessment of the Economic, Environmental and Social Impacts of NSW Agriculture's Wheat Breeding Program*, Economic Research Report No. 17, NSW Agriculture, Wagga Wagga.
- 18 Griffith, G.R., Davies, B.L., Alford, A.R., Herd, R.M., Parnell, P.F. and Hegarty, R.S. 2004, *An Assessment of the Economic, Environmental and Social Impacts of NSW Agriculture's Investment in the Net Feed Efficiency R,D&E Cluster*, Economic Research Report No. 18, NSW Department of Primary Industries, Armidale.
- 19 Scott, J.F. and Farquharson, R.J. 2004, *An Assessment of the Economic Impacts of NSW Agriculture's Research and Extension: Conservation Farming and Reduced Tillage in Northern NSW*, Economic Research Report No. 19, NSW Department of Primary Industries, Tamworth.
- 20 Scott, J.F., Farquharson, R.J. and Mullen, J.D. 2004, *Farming Systems in the Northern Cropping Region of NSW: An Economic Analysis*, Economic Research Report No. 20, NSW Department of Primary Industries, Tamworth.
- 21 Crean, J., Shaw, A., Singh, R. and Mullen, J.D. 2004, *An Assessment of the Economic, Environmental and Social Impacts of NSW Agriculture's Advisory Programs in Water Use Efficiency*, Economic Research Report No. 21, NSW Department of Primary Industries, Orange.
- 22 Mullen, J.D. 2004, *Evaluations in 2003 of Five Areas of Investment by NSW Agriculture: Summary*, Economic Research Report No. 22, NSW Department of Primary Industries, Orange.
- 23 Vere, D.T., Jones, R.E. and Dowling, P.M. 2004, *An Economic Evaluation of Research into the Improved Management of the Annual Grass Weed *Vulpia* in Temperate Pastures in South-Eastern Australia*, Economic Research Report No. 23, NSW Department of Primary Industries, Orange.
- 24 Jones, R.E. and Dowling, P.M. 2004, *Sustainability, Externalities and Economics: The Case of Temperate Perennial Grazing Systems in NSW*, Economic Research Report No. 24, NSW Department of Primary Industries, Orange.
- 25 Brennan, J.P. and Quade, K.J. 2004, *Analysis of the Impact of CIMMYT Research on the Australian Wheat Industry*, Economic Research Report No. 25, NSW Department of Primary Industries, Wagga Wagga.
- 26 Brennan, J.P., Sykes, J.D. and Scott, J.F. 2005, *Trends in Pulse and Oilseed Crops in Winter Cereal Rotations in NSW*, Economic Research Report No. 26, NSW Department of Primary Industries, Wagga Wagga.
- 27 Vere, D.T., Griffith, G.R. and Silvester, L. 2005, *Australian Sheep Industry CRC: Economic Evaluations of Scientific Research Programs*, Economic Research Report No. 27, NSW Department of Primary Industries, Orange.
- 28 Singh, R.P., Brennan, J.P., Lacy, J. and Steel, F. 2005, *An Assessment of the Economic, Environmental and Social Impacts of the Ricecheck Program*, Economic Research Report No. 28, NSW Department of Primary Industries, Yanco.
- 29 Jones, R., Saunders, G. and Balogh, S. 2005, *An Economic Evaluation of a Pest Management Control Program: "Outfox the Fox"*, Economic Research Report No. 29, NSW Department of Primary Industries, Orange.

- 30 Griffith, G.R., Parnell, P.F. and McKiernan, W. 2005, *The Economic, Environmental and Social Benefits to NSW from Investment in the CRC for Beef Genetics Technologies*, Economic Research Report No. 30, NSW Department of Primary Industries, Orange.
- 31 Khairo, S.A., Mullen, J.D., Hacker, R.B. & Patton, D.P 2006, *Farming Systems in the Western Division of NSW: An economic Analysis*, Economic Research Report No.31, NSW Department of Primary Industries, Trangie.
- 32 Griffith, G.R., Vere, D.T. and Jones, R.E. 2006, *Sheep CRC Renewal Proposal: Economic Evaluation of the Proposed Scientific Themes*, Economic Research Report No. 32, NSW Department of Primary Industries, Armidale, September.
- 33 Alford, A.R., Cafe, L.M., Greenwood, P.L. and Griffith, G.R. (2007), *The Economic Effects of Early-Life Nutritional Constraints in Crossbred Cattle Bred on the NSW North Coast*, Economic Research Report No. 33, NSW Department of Primary Industries, Armidale, February.